

Join us for a free NF1 PN educational virtual event.



Managing NF1 PN can be complex—that's why KOSELUGO® (selumetinib) comes with a team.

At Alexion, we know that support matters. That's why Koselugo comes with a team. Hear from a physician to learn more about neurofibromatosis type 1 with plexiform neurofibromas (NF1 PN).

At this event you will learn more about:

- Disease education from a physician
- Tips to prepare for treatment
- Info on Alexion support for NF1 families

For more information about Programs, reach out to a Patient Education Manager (PEM)

To find the PEM closest to you, use the PEM finder at **Alexiononesource.com/pem-finder**.

WHEN: Monday, August 7, 2023 7:00 PM ET WHERE: Virtual

To register and/or learn more about events, please scan the QR code, call 888-210-0174, or visit koselugo.com/events.



INDICATION

What is Koselugo?

Koselugo is a prescription medicine that is used to treat children 2 years of age and older with neurofibromatosis type 1 (NF1) who have plexiform neurofibromas that cannot be completely removed by surgery.

It is not known if Koselugo is safe and effective in children under 2 years of age.

IMPORTANT SAFETY INFORMATION

Before taking Koselugo, tell your healthcare provider about all of your medical conditions, including if you: • have heart problems

Please read additional Important Safety Information on the reverse side. Please see accompanying full

Prescribing Information , including Patient Information.

IMPORTANT SAFETY INFORMATION (continued)

Before taking Koselugo, tell your healthcare provider about all of your medical conditions, including if you:

- have eye problems
- are pregnant or plan to become pregnant. Koselugo can harm your unborn baby
 - Females who could become pregnant and males with female partners who could become pregnant should use effective birth control (contraception) during treatment with Koselugo and for 1 week after your last dose
 - Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Koselugo
- are breastfeeding or plan to breastfeed. It is not known if Koselugo passes into your breast milk
 - Do not breastfeed during treatment with Koselugo and for 1 week after your last dose
 - Talk to your healthcare provider about the best way to feed your baby during this time

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, or herbal supplements. Especially tell your healthcare provider if you are taking aspirin, blood thinners, or other medicines to treat blood clots. Koselugo contains vitamin E, which may increase your risk of bleeding.

Koselugo may cause serious side effects, including:

Heart problems. Koselugo can lower the amount of blood pumped by your heart, which is common and can also be severe. Your healthcare provider will do tests before and during treatment with Koselugo to check how well your heart is working. Tell your healthcare provider right away if you get any of the following signs or symptoms:

- persistent coughing or wheezing
 swelling of your ankles and feet
 increased heart rate
- shortness of breath
 tiredness

Eye problems. Koselugo can cause eye problems that can lead to blindness. Your healthcare provider will check your vision before and during treatment with Koselugo. Tell your healthcare provider right away if you get any of the following signs or symptoms:

blurred vision
 loss of vision
 dark spots in your vision
 other changes to your vision
 (floaters)

Severe diarrhea. Diarrhea is common with Koselugo and can also be severe. Tell your healthcare provider right away the first time that you get diarrhea during treatment with Koselugo. Your healthcare provider may give you medicine to help control your diarrhea and may tell you to drink more fluids.

Skin rash. Skin rashes are common with Koselugo and can also be severe. Tell your healthcare provider if you get any of the following signs or symptoms:

 rash that covers a large area of your body
 peeling skin
 blisters

Muscle problems (rhabdomyolysis). Muscle problems are common with Koselugo and can also be severe. Treatment with Koselugo may increase the level of a muscle enzyme in your blood, which may be a sign of muscle damage. Your healthcare provider should do a blood test to check your muscle enzyme levels before you start taking Koselugo and during treatment. Tell your healthcare provider right away if you get any of the following signs or symptoms:

- muscle aches or pain
- muscle spasms and weakness
- · dark, reddish urine

Your healthcare provider may change your dose, temporarily stop, or permanently ask you to stop taking Koselugo if you have any of these side effects.

The most common side effects of Koselugo are:

vomiting

 feeling of tiredness, weakness, or lacking energy

- stomach painnausea
- muscle and bone pain

dry skin

• fever

- inflammation of the mouth
- headache
- · redness around the fingernails
- itching

These are not all the possible side effects of Koselugo.

